

5.2 Simplex and Duality

The following linear programs form a primal dual pair:

$$z = \max\{c^T x \mid Ax = b, x \geq 0\}$$
$$w = \min\{b^T y \mid A^T y \geq c\}$$

This means for computing the dual of a standard form LP, we do not have non-negativity constraints for the dual variables.

Proof

Primal:

$$\begin{aligned} \max\{c^T x \mid Ax = b, x \geq 0\} \\ &= \max\{c^T x \mid Ax \leq b, -Ax \leq -b, x \geq 0\} \\ &= \max\{c^T x \mid \begin{bmatrix} A \\ -A \end{bmatrix} x \leq \begin{bmatrix} b \\ -b \end{bmatrix}, x \geq 0\} \end{aligned}$$

Dual:

$$\begin{aligned} \min\{[b^T \ -b^T]y \mid [A^T \ -A^T]y \geq c, y \geq 0\} \\ &= \min\left\{[b^T \ -b^T] \cdot \begin{bmatrix} y^+ \\ y^- \end{bmatrix} \mid [A^T \ -A^T] \cdot \begin{bmatrix} y^+ \\ y^- \end{bmatrix} \geq c, y^- \geq 0, y^+ \geq 0\right\} \\ &= \min\{b^T \cdot (y^+ - y^-) \mid A^T \cdot (y^+ - y^-) \geq c, y^- \geq 0, y^+ \geq 0\} \\ &= \min\{b^T y' \mid A^T y' \geq c\} \end{aligned}$$

Proof of Optimality Criterion for Simplex

Suppose that we have a basic feasible solution with **reduced cost**

$$\tilde{c} = c^T - c_B^T A_B^{-1} A \leq 0$$

This is equivalent to $A^T (A_B^{-1})^T c_B \geq c$

$y^* = (A_B^{-1})^T c_B$ is solution to the **dual** $\min\{b^T y \mid A^T y \geq c\}$.

$$\begin{aligned} b^T y^* &= (Ax^*)^T y^* = (A_B x_B^*)^T y^* \\ &= (A_B x_B^*)^T (A_B^{-1})^T c_B = (x_B^*)^T A_B^T (A_B^{-1})^T c_B \\ &= c^T x^* \end{aligned}$$

Hence, the solution is optimal.